


Study Title:
**Quantitative suspension test for evaluation of virucidal activity
in the medical area (Phase 2 Step1)**


Microbiological Solutions Limited (MSL)
Gollinrod, Walmersley, Bury, BL9 5NB, UK

Angela Davies, CEO

Customer: BCB International
Contact name: Benjamin Harrison
Email: bh@bcbin.com
Address: Howell House, Lamby Industrial Park, Wentloog Avenue, CF3 2EX'
PO/Quote number: Q002197/1



Megan Barrett
Laboratory Manager



Peter Thistlethwaite
Technical Projects Manager

The test results on this report refer only to the items tested as supplied by the customer. This report shall not be reproduced except in full and with written approval of Microbiological Solutions Ltd. All reports are archived for a minimum of 2 years. The sample will be retained for 1 month unless otherwise requested in writing.

Scope

The standard method BS EN 14476 describes a test method and the minimum requirements for virucidal activity of a chemical disinfectant and antiseptic products that form a homogenous physically stable preparation when diluted with hard water – or in the case of ready to use products that are not diluted when applied, - with water. Products can only be tested at a concentration of 80% (97% with a modified method for special cases) as some dilution is always produced by adding the test organisms and interfering substances. This European Standard applies to products that are used in the medical area in the fields of hygienic handrub, hygienic handwash, instrument disinfection by immersion, surface disinfection by wiping, spraying, flooding or other means and textile disinfection.

This European standard applies to areas and situations where disinfection is medically indicated. Such indication occurs in patient care, for example: In hospitals, in community medical facilities and in dental institutions or in clinics of schools, of kindergartens and of nursing homes, and may occur in the workplace and in the home. It may also include services such as laundries and kitchens supplying products directly for patients.

Outline of Test Method (Obligatory Test Conditions)

A sample of the test product is diluted in synthetic hard water in products diluted at point of use or water in the case of ready to use products is added to a test suspension of viruses in a solution of interfering substance. The mixture is maintained at one of the temperatures and contact times specified in the standard. At the end of this contact time, an aliquot is taken; the virucidal action in this portion is immediately suppressed by a validated method (dilutions of the sample in ice-cold cell maintenance medium). The dilutions are transferred into cell culture units either using monolayer or cell suspension. Infectivity tests are done either by plaque test or quantal tests. After incubation, the titres of infectivity are calculated according to Spearman and Käber or by plaque counting. Reduction of virus infectivity is calculated from differences of lg virus titres before (virus control) and after treatment with the product. The standard minimum spectrum of test organisms is Poliovirus, Adenovirus and Murine Norovirus.

Acceptance Criteria

The product when tested as above shall demonstrate at least a 4 log₁₀ reduction against the test virus. The test is deemed valid where all control requirements are met.

| Test information | | Deviation |
|--|--|-----------|
| Name of Product | Dr Brown's Hand Sanitiser | |
| Batch Number & Expiry Date | N/S | |
| Date of Delivery | 28/02/2020 | |
| Period of Analysis | 19/03/2020-23/03/2020 | |
| Manufacturer / Supplier | BCB International Ltd | |
| Storage Conditions | Ambient | |
| Appearance of the Product | Colourless gel | |
| Neutralisation Method | Dilution | |
| Product Diluent | Distilled water | |
| Test Concentrations | Neat (80%), Mid-range (50%) , Non active (0.1%) | |
| Experimental Conditions | Clean | |
| Interfering Substance | Clean 0.3g/l Bovine Albumin | |
| Test Temperature | 20°C ± 1°C | |
| Temperature of Incubation | 37°C ±1°C for 72hrs | |
| Identification of the Bacterial Strains: | Influenza H1N1 ATCC VR-1683 Feline Coronavirus, Strain Munich | 1 |
| Contact Times | 30 Seconds ± 10 s | |
| Stability and Appearance During Test | No Change Observed | |

Deviations from Standard Method

1 – The product was tested against nonstandard organisms Influenza H1N1 and Feline coronavirus, therefore reference inactivation controls were not performed due to no acceptance criteria available.

Test Result Summary

The test product received has achieved the following log reductions when tested under the condition stipulated in this report at a concentration of neat;


Influenza H1N1 - 3.25
Feline coronavirus – 4.46


See page 2 for acceptance criteria and raw data tables below for complete test results.


Summary Feline coronavirus

Microbiological Solutions Ltd
Gallinrod
Walmersley
Bury, BL9 5NB


Tel: 0844 824 6003
Email: info@rms.co
Web: www.rms.co
Company number: 4218514


| Controls | | | | | |
|---|---------------|--------------|------------|---------------|--------------------|
|  | | | | | |
| Conditions | Concentration | Contact time | log TCID50 | log reduction | Control validation |
| Virus control (water) | N/A | 30 seconds | 8.21 | N/A | Validated |
| Cytotoxicity (product) | Neat | N/A | 2.50 | N/A | Validated |
| Product supression control | Neat | Neat | 7.92 | 0.29 | Validated |


| Interference controls | | | | | |
|---|---------------|--------------|------------|----------------|--------------------|
|  | | | | | |
| Condition | Concentration | Contact time | log TCID50 | Log difference | Control validation |
| Interference control (untreated) | Neat | N/A | 9.00 | N/A | N/A |
| Interference control (treated) | Neat | N/A | 8.71 | 0.29 | Validated |

| Test Results | | | | |
|--|---------------|--------------|------------|---------------|
|  | | | | |
| Condition | Concentration | Contact time | log TCID50 | log reduction |
| Test product | Neat | 30 seconds | 3.75 | 4.46 |
| Test product | 50% | 30 seconds | 5.79 | 2.42 |
| Test product | 0.10% | 30 seconds | 7.92 | 0.29 |

Summary Influenza H1N1

| Controls | | | | | |
|---|---------------|--------------|------------|---------------|--------------------|
|  | | | | | |
| Conditions | Concentration | Contact time | log TCID50 | log reduction | Control validation |
| Virus control (water) | N/A | 30 Seconds | 7.08 | N/A | Validated |
| Cytotoxicity (product) | Neat | N/A | 2.50 | N/A | Validated |
| Product suppression control | Neat | Neat | 7.00 | 0.08 | Validated |

| Interference controls | | | | | |
|---|---------------|--------------|------------|----------------|--------------------|
|  | | | | | |
| Condition | Concentration | Contact time | log TCID50 | Log difference | Control validation |
| Interference control (untreated) | Neat | N/A | 9.00 | N/A | N/A |
| Interference control (treated) | Neat | N/A | 8.63 | 0.38 | Validated |

| Test Results | | | | |
|---|---------------|--------------|------------|---------------|
|  | | | | |
| Condition | Concentration | Contact time | log TCID50 | log reduction |
| Test product | Neat | 30 Seconds | 3.83 | 3.25 |
| Test product | 505% | 30 Seconds | 5.17 | 1.92 |
| Test product | 0.10% | 30 Seconds | 6.79 | 0.29 |

Raw data

| Virus control (water) | | Contact time | | | | | 30 seconds | | % CPE | p(1-p) |
|-----------------------|--------|--------------|---|---|---|---|------------|------------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -7 | 4 | 4 | 4 | 4 | 4 | 3 | 4 | 0.95833333 | 0.039931 | |
| -8 | 1 | 1 | 2 | 2 | 3 | 3 | 3 | 0.5 | 0.25 | |
| -9 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0.25 | 0.1875 | |

| Organism | | Feline Coronavirus | |
|----------|--|--------------------|--|
| | | Strain Munich | |
| d | | 1 | |
| sum px | | 2.71 | |
| n | | 8 | |
| SD50 | | -8.21 | |
| SE | | 0.26 | |
| kp | | -6 | |

| Cytotoxicity (product) | | Product concentration | | | | | Neat | | % CPE | p(1-p) |
|------------------------|--------|-----------------------|---|---|---|---|------|---|-------|--------|
| Dilution | Counts | | | | | | | | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |

| Organism | | Feline Coronavirus | |
|----------|--|--------------------|--|
| | | Strain Munich | |
| d | | 1 | |
| sum px | | 1.00 | |
| n | | 8 | |
| SD50 | | -2.50 | |
| SE | | 0.00 | |
| kp | | -2 | |

| Product supression control | | Product concentration | | | | | Neat | | % CPE | p(1-p) |
|----------------------------|--------|-----------------------|---|---|---|---|------|-----------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -7 | 3 | 3 | 3 | 4 | 4 | 4 | 4 | 0.875 | 0.109375 | |
| -8 | 2 | 2 | 2 | 1 | 1 | 1 | 1 | 0.375 | 0.234375 | |
| -9 | 1 | 1 | 0 | 1 | 1 | 1 | 0 | 0.1666667 | 0.138889 | |

| Organism | | Feline Coronavirus | |
|----------|--|--------------------|--|
| | | Strain Munich | |
| d | | 1 | |
| sum px | | 2.42 | |
| n | | 8 | |
| SD50 | | -7.92 | |
| SE | | 0.26 | |
| kp | | -6 | |

| Interference control (untreated) | | Product concentration | | | | | Neat | | % CPE | p(1-p) |
|----------------------------------|--------|-----------------------|---|---|---|---|------|------------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -1 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -7 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -8 | 3 | 3 | 4 | 4 | 4 | 4 | 4 | 0.91666667 | 0.076389 | |
| -9 | 2 | 2 | 2 | 1 | 1 | 1 | 1 | 0.375 | 0.234375 | |
| -10 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 0.20833333 | 0.164931 | |

| Organism | | Feline Coronavirus | |
|----------|--|--------------------|--|
| | | Strain Munich | |
| d | | 1 | |
| sum px | | 2.5 | |
| n | | 10 | |
| SD50 | | -9 | |
| SE | | 0.2299 | |
| kp | | -7 | |

| Interference control (treated) | | Product concentration | | | | | Neat | | % CPE | p(1-p) |
|--------------------------------|--------|-----------------------|---|---|---|---|------|------------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -1 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -7 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -8 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 0.75 | 0.1875 | |
| -9 | 2 | 2 | 2 | 1 | 1 | 1 | 0 | 0.33333333 | 0.222222 | |
| -10 | 1 | 1 | 0 | 0 | 0 | 1 | 1 | 0.125 | 0.109375 | |

| Organism | | Feline Coronavirus | |
|----------|--|--------------------|--|
| | | Strain Munich | |
| d | | 1 | |
| sum px | | 2.2083 | |
| n | | 10 | |
| SD50 | | -8.708 | |
| SE | | 0.2402 | |
| kp | | -7 | |

Raw data

| Test product | | Product concentration | | | | Neat | Contact time | | 30 seconds | |
|--------------|--------|-----------------------|---|---|---|------|--------------|--------|------------|--|
| Dilution | Counts | | | | | | % CPE | p(1-p) | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0.25 | 0.1875 | |
| -5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |

| Organism <i>Feline Coronavirus</i> | |
|------------------------------------|-------|
| Strain Munich | |
| d | 1 |
| sum px | 1.25 |
| n | 8 |
| SD50 | -3.75 |
| SE | 0.16 |
| xp | -3 |

| Test product | | Product concentration | | | | 50% | Contact time | | 30 seconds | |
|--------------|--------|-----------------------|---|---|---|-----|--------------|-------------|------------|--|
| Dilution | Counts | | | | | | % CPE | p(1-p) | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 3 | 3 | 0.91666667 | 0.076388889 | | |
| -6 | 2 | 2 | 2 | 1 | 1 | 0 | 0.33333333 | 0.22222222 | | |
| -7 | 1 | 0 | 0 | 0 | 0 | 0 | 0.04166667 | 0.039930556 | | |
| -8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |

| Organism <i>Feline Coronavirus</i> | |
|------------------------------------|-------|
| Strain Munich | |
| d | 1 |
| sum px | 2.29 |
| n | 8 |
| SD50 | -5.79 |
| SE | 0.22 |
| xp | -4 |

| Test product | | Product concentration | | | | 0.10% | Contact time | | 30 seconds | |
|--------------|--------|-----------------------|---|---|---|-------|--------------|-------------|------------|--|
| Dilution | Counts | | | | | | % CPE | p(1-p) | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -7 | 3 | 3 | 4 | 4 | 4 | 3 | 0.875 | 0.109375 | | |
| -8 | 2 | 2 | 1 | 1 | 2 | 2 | 0.41666667 | 0.243055556 | | |
| -9 | 1 | 1 | 1 | 0 | 0 | 0 | 0.125 | 0.109375 | | |

| Organism <i>Feline Coronavirus</i> | |
|------------------------------------|-------|
| Strain Munich | |
| d | 1 |
| sum px | 2.42 |
| n | 8 |
| SD50 | -7.92 |
| SE | 0.26 |
| xp | -6 |

Raw data

| Virus control (water) | | | Contact time | | | 30 Seconds | | | % CPE | p(1-p) |
|-----------------------|--------|---|--------------|---|---|------------|---|------------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 0.79166667 | 0.164931 | |
| -7 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 0.45833333 | 0.248264 | |
| -8 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0.25 | 0.1875 | |
| -9 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0.08333333 | 0.076389 | |

| Organism <i>Influenza</i> | |
|---------------------------|-------|
| H1N1 | |
| d | 1 |
| sum px | 2.58 |
| n | 8 |
| SD50 | -7.08 |
| SE | 0.31 |
| xp | -5 |

| Cytotoxicity (product) | | | Product concentration | | | Neat | | | % CPE | p(1-p) |
|------------------------|--------|---|-----------------------|---|---|------|---|---|-------|--------|
| Dilution | Counts | | | | | | | | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| -9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |

| Organism <i>Influenza</i> | |
|---------------------------|-------|
| H1N1 | |
| d | 1 |
| sum px | 1.00 |
| n | 8 |
| SD50 | -2.50 |
| SE | 0.00 |
| xp | -2 |

| Product suppression control | | | Product concentration | | | Neat | | | % CPE | p(1-p) |
|-----------------------------|--------|---|-----------------------|---|---|------|---|------------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 0.75 | 0.1875 | |
| -7 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 0.45833333 | 0.248264 | |
| -8 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0.25 | 0.1875 | |
| -9 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0.04166667 | 0.039931 | |

| Organism <i>Influenza</i> | |
|---------------------------|-------|
| H1N1 | |
| d | 1 |
| sum px | 2.50 |
| n | 8 |
| SD50 | -7.00 |
| SE | 0.31 |
| xp | -5 |

| Interference control (untreated) | | | Product concentration | | | Neat | | | % CPE | p(1-p) |
|----------------------------------|--------|---|-----------------------|---|---|------|---|------------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -1 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -7 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -8 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 0.79166667 | 0.164931 | |
| -9 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 0.58333333 | 0.243056 | |
| -10 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 0.125 | 0.109375 | |

| Organism <i>Influenza</i> | |
|---------------------------|--------|
| H1N1 | |
| d | 1 |
| sum px | 2.5 |
| n | 10 |
| SD50 | -9 |
| SE | 0.2398 |
| xp | -7 |

| Interference control (treated) | | | Product concentration | | | Neat | | | % CPE | p(1-p) |
|--------------------------------|--------|---|-----------------------|---|---|------|---|------------|----------|--------|
| Dilution | Counts | | | | | | | | | |
| -1 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -6 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -7 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | 0 | |
| -8 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 0.79166667 | 0.164931 | |
| -9 | 2 | 2 | 2 | 1 | 1 | 1 | 1 | 0.33333333 | 0.222222 | |
| -10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |

| Organism <i>Influenza</i> | |
|---------------------------|--------|
| H1N1 | |
| d | 1 |
| sum px | 2.125 |
| n | 10 |
| SD50 | -8.625 |
| SE | 0.2074 |
| xp | -7 |

Raw data

| Test product | Product concentration | | | | | Neat | Contact time | | 30 Seconds | |
|--------------|-----------------------|---|---|---|---|------|--------------|------------|------------|---|
| Dilution | Counts | | | | | | % CPE | p(1-p) | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -4 | 1 | 1 | 1 | 2 | 2 | 2 | 1 | 0.33333333 | 0.22222222 | 0 |
| -5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |
| -6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |
| -7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |
| -8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |
| -9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |

| Organism | | Influenza |
|----------|--|-----------|
| | | H1N1 |
| d | | 1 |
| sum px | | 1.33 |
| n | | 8 |
| SD50 | | -3.83 |
| SE | | 0.18 |
| xp | | -3 |

| Test product | Product concentration | | | | | 50% | Contact time | | 30 Seconds | |
|--------------|-----------------------|---|---|---|---|-----|--------------|------------|------------|---|
| Dilution | Counts | | | | | | % CPE | p(1-p) | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -5 | 3 | 3 | 3 | 2 | 2 | 2 | 1 | 0.58333333 | 0.24305556 | 0 |
| -6 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0.08333333 | 0.07638889 | 0 |
| -7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |
| -8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |
| -9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |

| Organism | | Influenza |
|----------|--|-----------|
| | | H1N1 |
| d | | 1 |
| sum px | | 1.67 |
| n | | 8 |
| SD50 | | -5.17 |
| SE | | 0.21 |
| xp | | -4 |

| Test product | Product concentration | | | | | 0.10% | Contact time | | 30 Seconds | |
|--------------|-----------------------|---|---|---|---|-------|--------------|------------|------------|---|
| Dilution | Counts | | | | | | % CPE | p(1-p) | | |
| -2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -5 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 1 | | 0 |
| -6 | 3 | 3 | 2 | 3 | 3 | 3 | 3 | 0.70833333 | 0.20659722 | 0 |
| -7 | 1 | 1 | 2 | 2 | 2 | 2 | 2 | 0.41666667 | 0.24305556 | 0 |
| -8 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0.16666667 | 0.13888889 | 0 |
| -9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 |

| Organism | | Influenza |
|----------|--|-----------|
| | | H1N1 |
| d | | 1 |
| sum px | | 2.29 |
| n | | 8 |
| SD50 | | -6.79 |
| SE | | 0.29 |
| xp | | -5 |

KEY

| | |
|-------------|--|
| CPE | Cytopathic effect |
| Counts | 0-4 indicating degree of cytopathic effect 0 = No effect, 1 = 25% CPE, 2 = 50% CPE, 3 = 75% CPE, 4 = 100% CPE |
| d | Dilution factor (log) |
| Sum px n | Sum of % CPE from the highest dilution showing 100% CPE to the lowest dilution assessed. Number of dilutions |
| SD50 | Dilution showing 50% of the end point according to Spearman-Kärber method |
| SE | Standard error |
| xp | Lowest dilution showing 100% CPE |
| TCID50 | Titre causing 50% of the end point according to Spearman-Kärber |
| PASS | = lg R greater than or equal to 4 |
| FAIL | = lg R less than 4 |
| > | greater than |
| < | less than |
| ≥ | equal to or greater than |
| ≤ | equal to or less than |

Calculation notes

In cases where the highest dilution assessed has not shown 100% CPE, the value has been calculated assuming the dilution above this would give 100% CPE and the corresponding value has been assigned as <x.

The standard requires the product suppression control to show a <0.5 log reduction in viral titre. In cases where the product has failed to achieve the required 4 log reduction, but the product suppression control shows a >0.5 log reduction the result has been deemed as valid for fail as the consequence of inadequate suppression would be a partially extended contact time which would generate false positives, but not false negatives.

A similar approach has been taken in regards to the cytotoxicity controls. The standard requires a 4-log difference between the cytotoxicity level and the viral titre. In cases where this is not obtained, but the log reduction observed by the product is within the difference between the cytotoxicity levels and the viral titre the result is deemed acceptable for a fail as there will be no impact on the determination of efficacy.